



## Botox, Dysport, Jeaveau, Xeomin Patient Instructions

### Pre Care Instructions

- Avoid the use of Aspirin, NSAIDS, Ginko Biloba, Garlic Supplements, Green Tea, Flax Oil, Cod Liver Oil, Vitamins A and E, and essential fatty acids for up to 1 week pre and post injection as they will increase your risk of bruising.
- Do not drink alcoholic beverages 24 hours before (or after) your treatment to avoid extra bruising
- You may take Arnica tablets 2-3 days prior to your injection to reduce the risk of bruising
- Do not apply products that are potentially irritating for 2 -3 days before and after treatment (examples- tretinoin/retin-a, glycolic acid, benzoyl peroxide, hydroquinone)
- You must inform the staff if you are pregnant or breastfeeding, are allergic to any of its ingredients in the above products, or have a history of neurologic disorders.

### Day of Treatment

- You may use topical anesthetic medication in office before your scheduled appointment (please arrive 15 minutes prior to procedure)
- Arrive with a clean face. Please do not wear makeup.
- You may experience a mild amount of tenderness or stinging during and following injection
- Redness and swelling are normal. Some bruising may also be visible.

### Post Care Instructions

- It is best to try to exercise your treated muscles for 1-2 hours after treatment (e.g. practice frowning, raise your eyebrows, and squint). This helps to work the product into your muscles.
- Ice (but avoid firm pressure) for 15 minutes 3 times the day of treatment if you have bruising
- You may take Acetaminophen/Tylenol if you experience any mild tenderness/discomfort
- Do not massage treated areas after treatment.
- After treatment there will be moderate swelling and redness with possibility of bruising. You may apply or take Arnica tablets to help decrease the amount of bruising.
- Avoid extended UV exposure until any redness/swelling has subsided. Apply SPF.
- Avoid strenuous exercise for 24 hours to reduce bruising. You may resume other normal activities/routines immediately.
- You may continue your topical products as mentioned above 3 days after injection.
- Avoid any laser or cosmetic treatments that massage or make the skin swell for 3 days.